






Astroparticle Physics

Lecture 2/2


Bradley J Kavanagh
Instituto de Fisica de Cantabria (CSIC-UC)

CERN Summer School - Tuesday 13th July 2021

The Sun



Supernovae




Quasars/AGN




Lecture 1

Lecture 2




????

Pulsars



Credit: Kevin Gill / Flickr

BH/NS Mergers



Credit: SXS Lensing

Credit: NASA/CXC/SAO

Credit: ESO/M. Kornmesser

Gravitational Waves


Einstein field equations of General Relativity:

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$$

Einstein tensor
(Gravity)

Stress-energy tensor
(Matter)

Space-time curvature specified by the metric, $g_{\mu\nu}$



Credit: ESA/C. Carreau

Linearise the field equations in vacuum:


$$g_{\mu\nu} \approx \eta_{\mu\nu} + h_{\mu\nu}$$

Wave-like solutions! **Gravitational Waves (GWs)**


$$\left(\frac{\partial^2}{\partial t^2} - \nabla^2 \right) h_{\mu\nu} = \square h_{\mu\nu} = 0$$

Indirect observation of GWs

GWs sourced by a time-dependent mass quadrupole moment.
E.g. compact object binaries...



Credit: Michael Kramer/Jodrell Bank Observatory/University of Manchester)




See e.g. [astro-ph/0407149](https://arxiv.org/abs/astro-ph/0407149)


Direct detection of GWs

www.ligo.caltech.edu/page/ligos-ifo

GW traveling into the screen causes (tiny) distortion:



[1708.00918](#)




Typical GW strain is $\Delta L/L \sim 10^{-23}$!


LIGO-Virgo-KAGRA (LVK)



Credit: LIGO, VIRGO & KAGRA.





GW frequency ~ twice orbital frequency.



[1906.03643](https://www.ligo.caltech.edu/page/strain-noise)

GW150914 - the first BH-BH merger


14 September 2015




Try it yourself! - <https://www.gw-openscience.org/tutorials/>

Properties of GW150914


Merger of two BHs - with masses $36 M_{\odot}$ and $29 M_{\odot}$ -
at a luminosity distance of $d_L \approx 200 - 600$ Mpc!




Combination $\mathcal{M} = (m_1 m_2)^{3/5} / (m_1 + m_2)^{1/5}$
(the chirp mass) is most tightly constrained.




Some degeneracy between distance and
binary orientation.



A whole catalog






A whole catalog

~ 50 GW merger events observed so far

[2007.07889](#)





[2106.15163](#)


GW170817 - the first NS-NS merger

17 August 2017 - observation of the merger
of two $\sim 1.5 - 2.0 M_{\odot}$ neutron stars

Localised to
within $\sim 30 \text{ deg}^2$



Multi-messenger follow-up




GW170817 merger occurred just two seconds before the gamma-ray burst GRB 170817A

Follow-up observations across the spectrum!

Sadly no neutrinos detected :(

What can we learn?

GW170817 resulted in a **kilonova**



[1710.05436](#)

Synthesis of r -process elements in neutron rich ejecta!

[1901.09044](#)

Extreme nuclear/quark physics!

[2103.16371](#)

Tests of general relativity!


[1710.06394](#)

Measurement of the Hubble Constant!


[1710.05835](#)

The Gravitational Wave Future

Planned Earth-based observatories such as Einstein Telescope:





Credit: Einstein Telescope



[1902.09485](#)

In addition, space-based detectors such as LISA will probe even lower frequencies (mHz) and therefore more massive systems (such as supermassive BH inspirals).

Dark Matter in Cosmology




Credit: ESA/Planck Collaboration


See [“Introduction to Cosmology” Lectures](#) by Paolo Creminelli

Dark Matter in Galaxies (1)

Both simulations and observations tell us: Galaxies contain lots of Dark Matter (DM)!



Illustris simulation - [1405.2921](#)




NB: Galaxy formation is messy and non-linear and still not fully understood


e.g. [1609.05917](#) vs [1610.07663](#)

Dark Matter in Galaxies (2)

Simulations point to Dark Matter halos with cuspy [NFW density profiles](#):



Aquarius simulation - [0809.0898](#)




DM density at Earth: $\rho_\chi \sim 5 \times 10^{-25} \text{ g/cm}^3$
 $\sim 0.3 \text{ GeV/cm}^3$
 $\sim 0.008 M_\odot/\text{pc}^3$

Dark Matter properties

Dark Matter must be:

- Non-baryonic
- Cold (i.e. slow-moving)
- (Almost) electrically neutral




*See additional slides...

Dark Matter properties

Dark Matter must be:


- Non-baryonic
- Cold (i.e. slow-moving)
- (Almost) electrically neutral



*See additional slides...

Dark Matter properties


- Dark Matter must be:
- Non-baryonic
 - Cold (i.e. slow-moving)
 - (Almost) electrically neutral



*See additional slides...


Direct detection of WIMPs on Earth

For WIMPs with GeV-scale masses, expect detectable nuclear recoils of energy $O(\text{keV})$



For sensible models, expect signal rates on the order of <1 event per kg per keV per day

No convincing signal yet!




Also possible to look for DM-electron scattering, depending on the model.

Indirect detection of Dark Matter

Look for signals of Dark Matter annihilation in regions of large DM density!

WIMP annihilation:




Annihilation cross section
(particle physics)

DM density distribution
(astrophysics)

$$\frac{d\Phi_\gamma}{dE_\gamma} = \frac{1}{4\pi} \frac{\langle \sigma_{\text{ann}} v \rangle}{2m_\chi^2} \frac{dN_\gamma}{dE_\gamma} \times \int_{d\Omega} d\Omega' \int_{los} \rho^2 dl (r, \theta')$$

Gamma-ray spectrum
(annihilation channel)



Aquarius simulation - [0809.0898](#)


Gamma-ray constraints

Fornax Dwarf Galaxy
(Satellite of the Milky Way)






Credit: ESO/Digitized Sky Survey 2

Fermi constraints from 15 Dwarf Spheroidal Galaxies:





Exact constraints depend on annihilation channel ($\chi\chi \rightarrow b\bar{b}$, $\chi\chi \rightarrow W^+W^-$, $\chi\chi \rightarrow e^+e^-$, etc.)

Galactic Centre Excess




Anti-proton excess (1)


Anti-protons are an excellent probe of New Physics - they're hard to make!



Anti-proton excess (2)



[1903.01472](#)





[1504.04276](#)

Several excesses point towards 60 GeV Dark Matter -
But modeling gamma-ray and cosmic-ray backgrounds is **hard**.

Point sources in the Galactic Centre

Galactic Centre excess could be due to a population of unresolved point sources (millisecond pulsars?)





Credit: Kevin Gill / Flickr

Credit: Christoph Weniger, UvA , © UvA/Princeton

High energy neutrinos

[1508.02500](#), [1712.07138](#)




Decays of super-heavy Dark Matter could contribute to the flux of PeV neutrinos:



Axion searches in the lab



Axions: light pseudoscalar particles

$$\begin{aligned}\mathcal{L} &\supset -\frac{1}{4}g_{a\gamma\gamma}aF_{\mu\nu}\tilde{F}^{\mu\nu} \\ &= -\frac{1}{4}g_{a\gamma\gamma}a\mathbf{E} \cdot \mathbf{B}\end{aligned}$$





Credit: Ciaran O'Hare, [AxionLimits](#)

Gamma-ray transparency and axions




[2012.15513](#)




[1302.1208](#)

The Sun



Supernovae





Quasars/AGN




Credit: NASA/CXC/SAO

Credit: ESO/M. Kornmesser




Pulsars



Credit: Kevin Gill / Flickr

BH/NS Mergers






Credit: SXS Lensing

Additional Slides



LISA - GWs in space!

Laser Interferometer Space Antenna
(Planned for the 2030s)






PBHs and Gravitational Waves



Gravitational Wave probes of DM



For more information about probing Dark Matter with Gravitational Waves, see [1907.10610](#)